

# Neural Networks: Optimization Part 1

Intro to Deep Learning, Spring 2023

#### Story so far

- Neural networks are universal approximators
  - Can model any odd thing
  - Provided they have the right architecture



- We must *train* them to approximate any function
  - Specify the architecture
  - Learn their weights and biases
- Networks are trained to minimize total "loss" on a training set
  - We do so through empirical risk minimization
- We use variants of gradient descent to do so
- The gradient of the error with respect to network parameters is computed through backpropagation

#### **Recap: Gradient Descent Algorithm**

- In order to minimize any function f(x) w.r.t. x
- Initialize:

$$-x^{0}$$

$$-k=0$$



• Do

$$-k = k + 1$$
  

$$-x^{k+1} = x^{k} - \eta \nabla_{x} f^{T}$$
  
while  $|f(x^{k}) - f(x^{k-1})| > \varepsilon$ 

#### Recap: Training Neural Nets by Gradient Descent

**Total training error:** 

$$Loss = \frac{1}{T} \sum_{t} Div(Y_t, d_t; W_1, W_2, \dots, W_K)$$

- Gradient descent algorithm:
- Initialize all weights **W**<sub>1</sub>, **W**<sub>2</sub>, ..., **W**<sub>K</sub>
- Do:
  - For every layer *k*, compute:
    - $\nabla_{\mathbf{W}_k} Loss = \frac{1}{T} \sum_t \nabla_{\mathbf{W}_k} Div(\mathbf{Y}_t, \mathbf{d}_t)$
    - $\mathbf{W}_k = \mathbf{W}_k \eta \nabla_{\mathbf{W}_k} Loss^T$
- Until Loss has converged

Computed using backprop

# Neural network training algorithm

- Initialize all weights and biases  $(\mathbf{W}_1, \mathbf{b}_1, \mathbf{W}_2, \mathbf{b}_2, \dots, \mathbf{W}_N, \mathbf{b}_N)$
- Do:
  - -Loss = 0
  - For all k, initialize  $\nabla_{\mathbf{W}_k} Loss = 0$ ,  $\nabla_{\mathbf{b}_k} Loss = 0$
  - For all t = 1:T # Loop through training instances
    - Forward pass : Compute
      - Output  $Y(X_t)$ ,
      - Divergence  $Div(Y_t, d_t)$
    - Backward pass: For all k compute:
      - $\nabla_{\mathbf{W}_k} Div(Y_t, d_t), \nabla_{\mathbf{b}_k} Div(Y_t, d_t)$
      - $\nabla_{\mathbf{W}_k} Loss += \nabla_{\mathbf{W}_k} Div(Y_t, d_t); \quad \nabla_{\mathbf{b}_k} Loss += \nabla_{\mathbf{b}_k} Div(Y_t, d_t)$
  - For all k, update:

 $\mathbf{W}_{k} = \mathbf{W}_{k} - \frac{\eta}{T} \left( \nabla_{\mathbf{W}_{k}} Loss \right)^{T}; \qquad \mathbf{b}_{k} = \mathbf{b}_{k} - \frac{\eta}{T} \left( \nabla_{\mathbf{W}_{k}} Loss \right)^{T}$ 

• Until *Loss* has converged

Computing gradient (uses backprop)

#### Issues

- Convergence: How well does it learn
   And how can we improve it
- How well will it generalize (outside training data)
- What does the output really mean?
- *Etc.*.

# Poll 0

Backpropagating from the kth layer, which is the derivative for the weights  $W_k$ ?

- $y_{k-1}$ .  $\nabla_{z_k} Div$ : The product of the output y of the k 1th layer and the derivative for the affine value z of the kth layer (in that order)
- $\nabla_{z_k} Div y_{k-1}$ : The product of the derivative for the affine value z at the kth layer and the output y of the k 1th layer (in that order)
- y<sup>T</sup><sub>k-1</sub>. ∇<sub>zk</sub> Div: The product of the transpose of the output y of the k –
   1th layer and the derivative for the affine value z of the kth layer (in that order)
- $\nabla_{z_k} Div. y_{k-1}^{\mathsf{T}}$ : The product of the derivative for the affine value z at the kth layer and the transpose output y of the k 1th layer (in that order)

# Poll 0

Backpropagating from the kth layer, which is the derivative for the weights  $W_k$ ?

- $y_{k-1}$ .  $\nabla_{z_k} Div$ : The product of the output y of the k 1th layer and the derivative for the affine value z of the kth layer (in that order)
- $\nabla_{z_k} Div y_{k-1}$ : The product of the derivative for the affine value z at the kth layer and the output y of the k 1th layer (in that order)
- y<sup>T</sup><sub>k-1</sub>. ∇<sub>zk</sub> Div: The product of the transpose of the output y of the k –
   1th layer and the derivative for the affine value z of the kth layer (in that order)
- $\nabla_{z_k} Div. y_{k-1}^{\mathsf{T}}$ : The product of the derivative for the affine value z at the kth layer and the transpose output y of the k 1th layer (in that order)

#### Onward



# Onward

- Does backprop always work?
- Convergence of gradient descent
  - Rates, restrictions,
  - Hessians
  - Acceleration and Nestorov
  - Alternate approaches
- Modifying the approach: Stochastic gradients
- Speedup extensions: RMSprop, Adagrad

# **Does backprop do the right thing?**

#### • Is backprop always right?

Assuming it actually finds the minimum of the divergence function?

(Actual question: Does gradient descent find the right solution, even when it finds the actual minimum)

#### **Recap: The differentiable activation**



- Threshold activation: Equivalent to counting errors
  - Shifting the threshold from T<sub>1</sub> to T<sub>2</sub> does not change classification error
  - Does not indicate if moving the threshold left was good or not



- Differentiable activation: Computes "distance to answer"
  - "Distance" == divergence
  - Perturbing the function changes this quantity,
    - Even if the classification error itself doesn't change

# Does backprop do the right thing?

#### • Is backprop always right?

- Assuming it actually finds the global minimum of the loss (average divergence)?
- In classification problems, the classification error is a non-differentiable function of weights
- The divergence function minimized is only a *proxy* for classification error
- Minimizing divergence may not minimize classification error



- Brady, Raghavan, Slawny, '89
- Simple problem, 3 training instances, single neuron
- Perceptron training rule trivially find a perfect solution

#### **Backprop vs. Perceptron**



- Back propagation using logistic function and  $L_2$ divergence  $(Div = (y - d)^2)$
- Unique minimum trivially proved to exist, backprop finds it

ν

Ζ



• From the three points we get three independent equations:

$$w_x \cdot 1 + w_y \cdot 0 + b = u$$
  
 $w_x \cdot 0 + w_y \cdot 1 + b = u$   
 $w_x \cdot -1 + w_y \cdot 0 + b = -u$ 

• Unique solution  $(w_x = u, w_x = u, b = 0)$  exists

•

represents a unique line regardless of the value of u

Y

Ζ



- Now add a fourth point
- t is very large (point near  $-\infty$ )
- Perceptron trivially finds a solution (may take t<sup>2</sup> iterations)

## Backprop



# Backprop

Notation:  

$$y = \sigma(z) = \text{logistic activation}$$
  
 $div_4 = (1 - \varepsilon - \sigma(-w_yt + b))^2$   
 $\frac{d \ div_4}{dw_y} = 2(1 - \varepsilon - \sigma(-w_yt + b))\sigma'(-w_yt + b)t$ 

• For very large positive t,  $|w_y| > \epsilon$  (where  $\mathbf{w} = [w_x, w_y, b]$ )

• 
$$(1 - \varepsilon - \sigma(-w_y t + b)) \to 1 \text{ as } t \to \infty$$

- $\sigma'(-w_yt+b) \to 0$  exponentially as  $t \to \infty$
- Therefore, for very large positive *t*

$$\frac{d \, div_4}{dw_y} = \frac{d \, div_4}{db} = 0$$



- The fourth point at (0, -t) does not change the gradient of the L<sub>2</sub> divergence near the optimal solution for 3 points
- The optimum solution for 3 points is also a broad *local* minimum (0 gradient) for the 4-point problem!
  - Will be found by backprop nearly all the time
    - Although the global minimum with unbounded weights will separate the classes correctly 20



- Local optimum solution found by backprop
- Does not separate the points even though the points are linearly separable!



- Solution found by backprop
- Does not separate the points even though the points are linearly separable!
- Compare to the perceptron: *Backpropagation fails to separate* where the perceptron succeeds

# **Backprop fails to separate where** perceptron succeeds γ

- Brady, Raghavan, Slawny, '89
- Several linearly separable training examples
- Simple setup: both backprop and perceptron algorithms find solutions



- Adding a "spoiler" (or a small number of sp
  - Perceptron finds the linear separator,



- Adding a "spoiler" (or a small number of spoilers)
  - Perceptron finds the linear separator,
  - Backprop does not find a separator
    - A single additional input does not change the loss function significantly
      - Assuming weights are constrained to be bounded

ν

X



- Adding a "spoiler" (or a small number of spoilers)
  - Perceptron finds the linear separator,
  - For bounded w, backprop does not find a separator
    - A single additional input does not change the loss function significantly



- Adding a "spoiler" (or a small number of spoilers)
  - Perceptron finds the linear separator,
  - For bounded w, backprop does not find a separator
    - A single additional input does not change the loss function significantly



- Adding a "spoiler" (or a small number of spoilers)
  - Perceptron finds the linear separator,
  - For bounded w, backprop does not find a separator
    - A single additional input does not change the loss function significantly

# So what is happening here?

- The perceptron may change greatly upon adding just a single new training instance
  - But it fits the training data well
  - The perceptron rule has *low bias*
    - Makes no errors if possible
  - But high variance
    - Swings wildly in response to small changes to input
- Backprop is minimally changed by new training instances
  - Prefers consistency over perfection
  - It is a *low-variance* estimator, at the potential cost of bias



- This is not restricted to single perceptrons
- An MLP learns non-linear decision boundaries that are determined from the entirety of the training data
- Adding a few "spoilers" will not change their behavior



- This is not restricted to single perceptrons
- An MLP learns non-linear decision boundaries that are determined from the entirety of the training data
- Adding a few "spoilers" will not change their behavior

#### **Backpropagation: Finding the separator**

- Backpropagation will often not find a separating solution *even though the solution is within the class of functions learnable by the network*
- This is because the separating solution is not a feasible optimum for the loss function
- One resulting benefit is that a backprop-trained neural network classifier has lower variance than an optimal classifier for the training data

#### Poll 1

Minimizing the (differentiable) loss function will also minimize classification error, true or false

- True
- False

#### Poll 1

Minimizing the (differentiable) loss function will also minimize classification error, true or false

- True
- False (true)

## **The Loss Surface**

- The example (and statements) earlier assumed the loss
   objective had a single global
   optimum that could be found
  - Statement about variance is assuming global optimum
- What about local optima



#### **The Loss Surface**

- Popular hypothesis:
  - In large networks, saddle points are far more common than local minima
    - Frequency of occurrence exponential in network size
  - Most local minima are equivalent
    - And close to global minimum
  - This is not true for small networks
- Saddle point: A point where
  - The slope is zero
  - The surface increases in some directions, but decreases in others
    - Some of the Eigenvalues of the Hessian are positive; others are negative
  - Gradient descent algorithms often get "stuck" in saddle points




# **The Controversial Loss Surface**

- Baldi and Hornik (89), "Neural Networks and Principal Component Analysis: Learning from Examples Without Local Minima" : An MLP with a single hidden layer has only saddle points and no local Minima
- **Dauphin et. al (2015),** *"Identifying and attacking the saddle point problem in high-dimensional non-convex optimization"* : An exponential number of saddle points in large networks
- Chomoranksa et. al (2015), "The loss surface of multilayer networks" : For large networks, most local minima lie in a band and are equivalent
  - Based on analysis of spin glass models
- Swirscz et. al. (2016), "Local minima in training of deep networks", In networks of finite size, trained on finite data, you *can* have horrible local minima
- Watch this space...

## Story so far

- Neural nets can be trained via gradient descent that minimizes a loss function
- Backpropagation can be used to derive the derivatives of the loss
- Backprop *is not guaranteed* to find a "true" solution, even if it exists, and lies within the capacity of the network to model
  - The optimum for the loss function may not be the "true" solution
- For large networks, the loss function may have a large number of unpleasant saddle points or local minima
  - Which backpropagation may find

#### Convergence

- In the discussion so far we have assumed the training arrives at a local minimum
- Does it always converge?
- How long does it take?
- Hard to analyze for an MLP, but we can look at the problem through the lens of convex optimization

#### A quick tour of (convex) optimization



"I'm searching for my keys."

# **Convex Loss Functions**

- A surface is "convex" if it is continuously curving upward
  - We can connect any two points on or above the surface without intersecting it
  - Many mathematical definitions that are equivalent
- Caveat: Neural network loss surface is generally not convex
  - Streetlight effect





# **Convergence of gradient descent**

- An iterative algorithm is said to converge to a solution if the value updates arrive at a fixed point
  - Where the gradient is 0 and further updates do not change the estimate
- The algorithm may not actually converge
  - It may jitter around the local minimum
  - It may even diverge
- Conditions for convergence?







#### **Convergence and convergence rate**

- Convergence rate: How fast the iterations arrive at the solution
- Generally quantified as

$$R = \frac{\left| f(x^{(k+1)}) - f(x^*) \right|}{\left| f(x^{(k)}) - f(x^*) \right|}$$

- $x^{(k+1)}$  is the k-th iteration
- $x^*$  is the optimal value of x
- If *R* is a constant (or upper bounded), the convergence is *linear* 
  - In reality, its arriving at the solution exponentially fast

$$|f(x^{(k)}) - f(x^*)| \le R^k |f(x^{(0)}) - f(x^*)|$$



## **Convergence for quadratic surfaces**

 $Minimize E = \frac{1}{2}aw^2 + bw + c$ 

 $\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta \frac{dE(\mathbf{w}^{(k)})}{d\mathbf{w}}$ 

Gradient descent with fixed step size  $\eta$  to estimate scalar parameter w

- $E(\omega)$ (f)a)  $W^{(k)}$
- Gradient descent to find the optimum of a quadratic, starting from w<sup>(k)</sup>
- Assuming fixed step size  $\eta$
- What is the optimal step size
  η to get there fastest?

## **Convergence for quadratic surfaces**

$$E = \frac{1}{2}aw^2 + bw + c$$
$$w^{(k+1)} = w^{(k)} - \eta \frac{dE(w^{(k)})}{dw}$$

- Any quadratic objective can be written as  $E(w) = E(w^{(k)}) + E'(w^{(k)})(w - w^{(k)}) + \frac{1}{2}E''(w^{(k)})(w - w^{(k)})^2$ 
  - Taylor expansion



• Minimizing w.r.t w, we get (Newton's method)  $w_{min} = w^{(k)} - E'' (w^{(k)})^{-1} E' (w^{(k)})$ 

• Note:

ullet

$$\frac{dE(\mathbf{w}^{(k)})}{d\mathbf{w}} = E'(\mathbf{w}^{(k)})$$

• Comparing to the gradient descent rule, we see that we can arrive at the optimum in a single step using the optimum step size

$$\eta_{opt} = E'' \left( \mathbf{w}^{(k)} \right)^{-1} = \boldsymbol{a}^{-1}$$

# With non-optimal step size

$$w^{(k+1)} = w^{(k)} - \eta \frac{dE(w^{(k)})}{dw}$$





Gradient descent with fixed step size  $\eta$  to estimate scalar parameter w

- For  $\eta < \eta_{opt}$  the algorithm will converge monotonically
- For  $2\eta_{opt} > \eta > \eta_{opt}$  we have oscillating convergence
- For  $\eta > 2\eta_{opt}$  we get divergence

# For generic differentiable convex objectives



• Any differentiable convex objective E(w) can be approximated as

$$E \approx E(\mathbf{w}^{(k)}) + (w - \mathbf{w}^{(k)}) \frac{dE(\mathbf{w}^{(k)})}{dw} + \frac{1}{2}(w - \mathbf{w}^{(k)})^2 \frac{d^2E(\mathbf{w}^{(k)})}{dw^2} + \cdots$$

Taylor expansion

• Using the same logic as before, we get (Newton's method)

$$\eta_{opt} = \left(\frac{d^2 E(\mathbf{w}^{(k)})}{dw^2}\right)^{-1}$$

• We can get divergence if  $\eta \ge 2\eta_{opt}$ 

#### For functions of *multivariate* inputs

 $E = g(\mathbf{w}), \mathbf{w}$  is a vector  $\mathbf{w} = [w_1, w_2, \dots, w_N]$ 

• Consider a simple quadratic convex (paraboloid) function

$$E = \frac{1}{2}\mathbf{w}^T \mathbf{A}\mathbf{w} + \mathbf{w}^T \mathbf{b} + c$$

- Since  $E^T = E$  (*E* is scalar), **A** can always be made symmetric
  - For strictly **convex** *E*, **A** is always positive definite, and has positive eigenvalues
- When **A** is diagonal:

$$E = \frac{1}{2} \sum_{i} (a_{ii} w_i^2 + b_i w_i) + c$$

- The  $w_i$ s are *uncoupled*
- For paraboloid (*convex*) E, the  $a_{ii}$  values are all positive
- Just a sum of N independent quadratic functions

#### **Multivariate Quadratic with Diagonal A**



• Equal-value contours will ellipses with principal axes parallel to the spatial axes

#### **Multivariate Quadratic with Diagonal A**



- Equal-value contours will be parallel to the axes
  - All "slices" parallel to an axis are shifted versions of one another

$$E = \frac{1}{2}a_{ii}w_i^2 + b_iw_i + c + C(\neg w_i)$$

#### **Multivariate Quadratic with Diagonal A**



- Equal-value contours will be parallel to the axis
  - All "slices" parallel to an axis are shifted versions of one another

$$E = \frac{1}{2}a_{ii}w_i^2 + b_iw_i + c + C(\neg w_i)$$

#### "Descents" are uncoupled



- The optimum of each coordinate is not affected by the other coordinates
  I.e. we could optimize each coordinate independently
- Note: Optimal learning rate is different for the different coordinates

## Vector update rule



$$\mathbf{w}^{(k+1)} \leftarrow \mathbf{w}^{(k)} - \eta \nabla_{\mathbf{w}} E^{\top}$$

$$w_i^{(k+1)} = w_i^{(k)} - \eta \frac{\partial E\left(w_i^{(k)}\right)}{d\partial w}$$

- Conventional vector update rules for gradient descent: update entire vector against direction of gradient
  - Note : Gradient is perpendicular to equal value contour
  - The same learning rate is applied to all components

## **Problem with vector update rule**

$$\mathbf{w}^{(k+1)} \leftarrow \mathbf{w}^{(k)} - \eta \nabla_{\mathbf{w}} E^{T}$$
$$w_{i}^{(k+1)} = w_{i}^{(k)} - \eta \frac{\partial E\left(w_{i}^{(k)}\right)}{\partial w}$$
$$\eta_{i,opt} = \left(\frac{\partial^{2} E\left(w_{i}^{(k)}\right)}{\partial w_{i}^{2}}\right)^{-1} = a_{ii}^{-1}$$

#### **Dependence on learning rate**

10

10

20

20





- $\eta_{1,opt} = 1; \ \eta_{2,opt} = 0.33$
- $\eta = 2.1\eta_{2,opt}$
- $\eta = 2\eta_{2,opt}$
- $\eta = 1.5\eta_{2,opt}$
- $\eta = \eta_{2,opt}$
- $\eta = 0.75\eta_{2,opt}$

# **Problem with vector update rule**

$$\mathbf{w}^{(k+1)} \leftarrow \mathbf{w}^{(k)} - \eta \nabla_{\mathbf{w}} E^{T} \qquad w_{i}^{(k+1)} = w_{i}^{(k)} - \eta \frac{\partial E\left(w_{i}^{(k)}\right)}{\partial w}$$
$$\eta_{i,opt} = \left(\frac{\partial^{2} E\left(w_{i}^{(k)}\right)}{\partial w_{i}^{2}}\right)^{-1} = a_{ii}^{-1}$$

• The learning rate must be lower than twice the *smallest* optimal learning rate for any component

 $\eta < 2 \min_{i} \eta_{i,opt}$ 

Otherwise the learning will diverge

• This, however, makes the learning very slow

- And will oscillate in all directions where  $\eta_{i,opt} \leq \eta < 2\eta_{i,opt}$ 

#### **Dependence on learning rate**



•  $\eta_{1,opt} = 1; \ \eta_{2,opt} = 0.91; \qquad \eta = 1.9 \ \eta_{2,opt}$ 

# Generic differentiable *multivariate* convex functions



- For generic convex multivariate functions (not necessarily quadratic), we can employ quadratic Taylor series expansions and much of the analysis still applies
- Taylor expansion

$$E(\mathbf{w}) \approx E(\mathbf{w}^{(k)}) + \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)}) (\mathbf{w} - \mathbf{w}^{(k)}) + \frac{1}{2} (\mathbf{w} - \mathbf{w}^{(k)})^T H_E(\mathbf{w}^{(k)}) (\mathbf{w} - \mathbf{w}^{(k)})$$

- The optimal step size is inversely proportional to the Eigen values of the Hessian
  - The second derivative along the orthogonal coordinates
  - For the smoothest convergence, these must all be equal

#### Convergence

- Convergence behaviors become increasingly unpredictable as dimensions increase
- For the fastest convergence, ideally, the learning rate  $\eta$  must be close to both, the largest  $\eta_{i,opt}$  and the smallest  $\eta_{i,opt}$ 
  - To ensure convergence in every direction
  - Generally infeasible
- Convergence is particularly slow if  $\frac{\max_{i} \eta_{i,opt}}{\min_{i} \eta_{i,opt}}$  is large
  - The "condition" number
    - Must be close to 1.0 for fast convergence
- Following (hidden) slides discuss solutions that "normalize the space by stretching different directions differently to standardize optimal step size
  - A big topic for optimization
  - Unfortunately, infeasible for neural networks

# **Comments on the quadratic**

- Why are we talking about quadratics?
  - Quadratic functions form some kind of benchmark
  - Convergence of gradient descent is linear
    - Meaning it converges to solution exponentially fast
- The convergence for other kinds of functions can be viewed against this benchmark
- Actual losses will not be quadratic, but may locally have other structure
  - Local between current location and nearest local minimum
- Some examples in the following slides..
  - Strong convexity
  - Lifschitz continuity
  - Lifschitz smoothness
  - ..and how they affect convergence of gradient descent

#### **Quadratic convexity**



- A quadratic function has the form  $\frac{1}{2}\mathbf{w}^T\mathbf{A}\mathbf{w} + \mathbf{w}^T\mathbf{b} + c$ 
  - Every "slice" is a quadratic bowl
- In some sense, the "standard" for gradient-descent based optimization
  - Others convex functions will be steeper in some regions, but flatter in others
- Gradient descent solution will have linear convergence
  - Take  $O(\log 1/\varepsilon)$  steps to get within  $\varepsilon$  of the optimal solution

# **Strong convexity**



- A strongly convex function is *at least* quadratic in its convexity
  - Has a lower bound to its second derivative
- The function sits within a quadratic bowl
  - At any location, you can draw a quadratic bowl of fixed convexity (quadratic constant equal to lower bound of 2<sup>nd</sup> derivative) touching the function at that point, which contains it
- Convergence of gradient descent algorithms at least as good as that of the enclosing quadratic

## **Strong convexity**



- A strongly convex function is *at least* quadratic in its convexity
  - Has a lower bound to its second derivative
- The function sits within a quadratic bowl
  - At any location, you can draw a quadratic bowl of fixed convexity (quadratic constant equal to lower bound of 2<sup>nd</sup> derivative) touching the function at that point, which contains it
- Convergence of gradient descent algorithms at least as good as that of the enclosing quadratic

# **Types of continuity**



- Most functions are not strongly convex (if they are convex)
- Instead we will talk in terms of Lifschitz smoothness
- But first : a definition
- *Lifschitz continuous*: The function always lies outside a cone
  - The slope of the outer surface is the Lifschitz constant

$$-|f(x) - f(y)| \le L|x - y|$$

## Lifschitz smoothness



- Lifschitz smooth: The function's *derivative* is Lifschitz continuous
  - Need not be convex (or even differentiable)
  - Has an upper bound on second derivative (if it exists)
- Can always place a quadratic bowl of a fixed curvature within the function
  - Minimum curvature of quadratic must be >= upper bound of second derivative of function (if it exists)

## Lifschitz smoothness



- Lifschitz smooth: The function's *derivative* is Lifschitz continuous
  - Need not be convex (or even differentiable)
  - Has an *upper bound* on second derivative (if it exists)
- Can always place a quadratic bowl of a fixed curvature within the function
  - Minimum curvature of quadratic must be >= upper bound of second derivative of function (if it exists)

# **Types of smoothness**



- A function can be both strongly convex and Lipschitz smooth
  - Second derivative has upper and lower bounds
  - Convergence depends on curvature of strong convexity (at least linear)
- A function can be convex and Lifschitz smooth, but not strongly convex
  - Convex, but upper bound on second derivative
  - Weaker convergence guarantees, if any (at best linear)
  - This is often a reasonable assumption for the local structure of your loss function

# **Types of smoothness**



- A function can be both strongly convex and Lipschitz smooth
  - Second derivative has upper and lower bounds
  - Convergence depends on curvature of strong convexity (at least linear)
- A function can be convex and Lifschitz smooth, but not strongly convex
  - Convex, but upper bound on second derivative
  - Weaker convergence guarantees, if any (at best linear)
  - This is often a reasonable assumption for the local structure of your loss function

# **Convergence Problems**

- For quadratic (strongly) convex functions, gradient descent is exponentially fast
  - Linear convergence
    - Assuming learning rate is non-divergent
- For generic (Lifschitz Smooth) convex functions however, it is very slow

$$|f(w^{(k)}) - f(w^*)| \propto \frac{1}{k} |f(w^{(0)}) - f(w^*)|$$

And inversely proportional to learning rate

$$\left|f(w^{(k)}) - f(w^*)\right| \le \frac{1}{2\eta k} \left|w^{(0)} - w^*\right|$$

- Takes  $O(1/\epsilon)$  iterations to get to within  $\epsilon$  of the solution
- An inappropriate learning rate will destroy your happiness
- Second order methods will *locally* convert the loss function to quadratic
  - Convergence behavior will still depend on the nature of the original function
- Continuing with the quadratic-based explanation...

## Convergence

- Convergence behaviors become increasingly unpredictable as dimensions increase
- For the fastest convergence, ideally, the learning rate  $\eta$ must be close to both, the largest  $\eta_{i,opt}$  and the smallest  $\eta_{i,opt}$ 
  - To ensure convergence in every direction
  - Generally infeasible
- Convergence is particularly slow if  $\frac{\max_{i} \eta_{i,opt}}{\min_{i} \eta_{i,opt}}$  is large

The "condition" number is small



- The objective function has different eccentricities in different directions
  - Resulting in different optimal learning rates for different directions
  - The problem is more difficult when the ellipsoid is not axis aligned: the steps along the two directions are coupled! Moving in one direction changes the gradient along the other
- Solution: *Normalize* the objective to have identical eccentricity in all directions
  - Then all of them will have identical optimal learning rates
  - Easier to find a working learning rate

## **Solution: Scale the axes**



- Scale (and rotate) the axes, such that all of them have identical (identity) "spread"
  - Equal-value contours are circular
  - Movement along the coordinate axes become independent
- Note: equation of a quadratic surface with circular equal-value contours can be written as

$$E = \frac{1}{2}\widehat{\mathbf{w}}^T\widehat{\mathbf{w}} + \widehat{\mathbf{b}}^T\widehat{\mathbf{w}} + c$$
• Original equation:

$$E = \frac{1}{2}\mathbf{w}^T \mathbf{A}\mathbf{w} + \mathbf{b}^T \mathbf{w} + c$$

• We want to find a (diagonal) scaling matrix *S* such that

$$\mathbf{S} = \begin{bmatrix} s_1 & \cdots & 0\\ \vdots & \ddots & \vdots\\ 0 & \cdots & s_N \end{bmatrix}, \qquad \widehat{\mathbf{w}} = \mathbf{S}\mathbf{w}$$

• And

$$E = \frac{1}{2}\widehat{\mathbf{w}}^T\widehat{\mathbf{w}} + \widehat{\mathbf{b}}^T\widehat{\mathbf{w}} + c$$

• Original equation:

$$E = \frac{1}{2}\mathbf{w}^T \mathbf{A}\mathbf{w} + \mathbf{b}^T \mathbf{w} + c$$

• We want to find a (diagonal) scaling matrix S such that

$$\mathbf{S} = \begin{bmatrix} s_1 & \cdots & 0\\ \vdots & \ddots & \vdots\\ 0 & \cdots & s_N \end{bmatrix}, \qquad \widehat{\mathbf{w}} = \mathbf{S}\mathbf{w}$$

• And

$$E = \frac{1}{2}\widehat{\mathbf{w}}^T\widehat{\mathbf{w}} + \widehat{\mathbf{b}}^T\widehat{\mathbf{w}} + c$$

By inspection:  $S = A^{0.5}$ 

• We have

$$E = \frac{1}{2} \mathbf{w}^T \mathbf{A} \mathbf{w} + \mathbf{b}^T \mathbf{w} + c$$
$$\widehat{\mathbf{w}} = S\mathbf{w}$$
$$E = \frac{1}{2} \widehat{\mathbf{w}}^T \widehat{\mathbf{w}} + \widehat{\mathbf{b}}^T \widehat{\mathbf{w}} + c$$
$$= \frac{1}{2} \mathbf{w}^T S^T S\mathbf{w} + \widehat{\mathbf{b}}^T S\mathbf{w} + c$$

• Equating linear and quadratic coefficients, we get

• Solving: 
$$S = A^{0.5}$$
,  $\hat{b} = A^{-0.5}b$ 

• We have

$$E = \frac{1}{2} \mathbf{w}^T \mathbf{A} \mathbf{w} + \mathbf{b}^T \mathbf{w} + c$$
$$\widehat{\mathbf{w}} = \mathbf{S} \mathbf{w}$$
$$E = \frac{1}{2} \widehat{\mathbf{w}}^T \widehat{\mathbf{w}} + \widehat{\mathbf{b}}^T \widehat{\mathbf{w}} + c$$

• Solving for <mark>S</mark> we get

$$\widehat{\mathbf{w}} = \mathbf{A}^{0.5} \mathbf{w}, \qquad \widehat{\mathbf{b}} = \mathbf{A}^{-0.5} \mathbf{b}$$

• We have

$$E = \frac{1}{2}\mathbf{w}^{T}\mathbf{A}\mathbf{w} + \mathbf{b}^{T}\mathbf{w} + c$$
$$\widehat{\mathbf{w}} = S\mathbf{w}$$
$$E = \frac{1}{2}\widehat{\mathbf{w}}^{T}\widehat{\mathbf{w}} + \widehat{\mathbf{b}}^{T}\widehat{\mathbf{w}} + c$$

• Solving for <mark>S</mark> we get

$$\widehat{\mathbf{w}} = \widehat{\mathbf{A}^{0.5}}\mathbf{w},$$

$$\hat{\mathbf{b}} = \mathbf{A}^{-0.5}\mathbf{b}$$

## The Inverse Square Root of A

- For *any* positive definite **A**, we can write  $\mathbf{A} = \mathbf{E} \mathbf{\Lambda} \mathbf{E}^{\mathrm{T}}$ 
  - Eigen decomposition
  - E is an orthogonal matrix
  - $-\Lambda$  is a diagonal matrix of non-zero diagonal entries
- Defining  $\mathbf{A}^{0.5} = \mathbf{E} \mathbf{\Lambda}^{0.5} \mathbf{E}^{\mathrm{T}}$ - Check  $(\mathbf{A}^{0.5})^{\mathrm{T}} \mathbf{A}^{0.5} = \mathbf{E} \mathbf{\Lambda} \mathbf{E}^{\mathrm{T}} = \mathbf{A}$
- Defining  $\mathbf{A}^{-0.5} = \mathbf{E} \mathbf{\Lambda}^{-0.5} \mathbf{E}^{\mathrm{T}}$ - Check:  $(\mathbf{A}^{-0.5})^{\mathrm{T}} \mathbf{A}^{-0.5} = \mathbf{E} \mathbf{\Lambda}^{-1} \mathbf{E}^{\mathrm{T}} = \mathbf{A}^{-1}$

## **Returning to our problem**



$$E = \frac{1}{2}\widehat{\mathbf{w}}^T\widehat{\mathbf{w}} + \widehat{\mathbf{b}}^T\widehat{\mathbf{w}} + c$$

• Computing the gradient, and noting that  $A^{0.5}$  is symmetric, we can relate  $\nabla_{\widehat{w}}E$  and  $\nabla_{w}E$ :

$$\nabla_{\widehat{\mathbf{w}}} E = \widehat{\mathbf{w}}^T + \widehat{\mathbf{b}}^T$$
$$= \mathbf{w}^T \mathbf{A}^{0.5} + \mathbf{b}^T \mathbf{A}^{-0.5}$$
$$= (\mathbf{w}^T \mathbf{A} + \mathbf{b}^T) \mathbf{A}^{-0.5}$$
$$= \nabla_{\mathbf{w}} E \cdot \mathbf{A}^{-0.5}$$

#### **Returning to our problem**



 $E = \frac{1}{2}\widehat{\mathbf{w}}^T\widehat{\mathbf{w}} + \widehat{\mathbf{b}}^T\widehat{\mathbf{w}} + c$ 

Gradient descent rule:

$$-\widehat{\mathbf{w}}^{(k+1)} = \widehat{\mathbf{w}}^{(k)} - \eta \nabla_{\widehat{\mathbf{w}}} E(\widehat{\mathbf{w}}^{(k)})^{T}$$

Learning rate is now independent of direction

• Using  $\widehat{\mathbf{w}} = \mathbf{A}^{0.5} \mathbf{w}$ , and  $\nabla_{\widehat{\mathbf{w}}} E(\widehat{\mathbf{w}})^T = \mathbf{A}^{-0.5} \nabla_{\mathbf{w}} E(\mathbf{w})^T$ 

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta \mathbf{A}^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$

### **Modified update rule**



• 
$$\widehat{\mathbf{w}}^{(k+1)} = \widehat{\mathbf{w}}^{(k)} - \eta \nabla_{\widehat{\mathbf{w}}} E(\widehat{\mathbf{w}}^{(k)})^T$$

Leads to the modified gradient descent rule

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta \mathbf{A}^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$

## For non-axis-aligned quadratics..



 $E = \frac{1}{2}\mathbf{w}^T \mathbf{A}\mathbf{w} + \mathbf{w}^T \mathbf{b} + c$ 

$$E = \frac{1}{2} \sum_{i} a_{ii} w_i^2 + \sum_{i \neq j} a_{ij} w_i w_j$$
$$+ \sum_{i} b_i w_i + c$$

- If A is not diagonal, the contours are not axis-aligned
  - Because of the cross-terms  $a_{ij}w_iw_j$
  - The major axes of the ellipsoids are the *Eigenvectors* of **A**, and their diameters are proportional to the Eigen values of **A**
- But this does not affect the discussion
  - This is merely a rotation of the space from the axis-aligned case
  - The component-wise optimal learning rates along the major and minor axes of the equalcontour ellipsoids will be different, causing problems
    - The optimal rates along the axes are Inversely proportional to the eigenvalues of A



- The component-wise optimal learning rates along the major and minor axes of the contour ellipsoids will differ, causing problems
  - Inversely proportional to the *eigenvalues* of A
- This can be fixed as before by rotating and resizing the different directions to obtain the same *normalized* update rule as before:  $\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta \mathbf{A}^{-1} \mathbf{b}$

## Generic differentiable *multivariate* convex functions



Taylor expansion

 $E(\mathbf{w}) \approx E(\mathbf{w}^{(k)}) + \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)}) (\mathbf{w} - \mathbf{w}^{(k)}) + \frac{1}{2} (\mathbf{w} - \mathbf{w}^{(k)})^{T} H_{E}(\mathbf{w}^{(k)}) (\mathbf{w} - \mathbf{w}^{(k)}) + \cdots$ 



#### Generic differentiable *multivariate* convex functions



• Taylor expansion

 $E(\mathbf{w}) \approx E(\mathbf{w}^{(k)}) + \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)}) (\mathbf{w} - \mathbf{w}^{(k)}) + \frac{1}{2} (\mathbf{w} - \mathbf{w}^{(k)})^{T} H_{E}(\mathbf{w}^{(k)}) (\mathbf{w} - \mathbf{w}^{(k)}) + \cdots$ 

- Note that this has the form  $\frac{1}{2}\mathbf{w}^T\mathbf{A}\mathbf{w} + \mathbf{w}^T\mathbf{b} + c$
- Using the same logic as before, we get the normalized update rule

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E (\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E (\mathbf{w}^{(k)})^T$$

- For a quadratic function, the optimal  $\eta$  is 1 (which is exactly Newton's method)
  - And should not be greater than 2!



Fit a quadratic at each point and find the minimum of that quadratic

Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^T$$

$$-\eta = 1$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$

$$-\eta = 1$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^T$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^T$$

$$-\eta = 1$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$

$$-\eta = 1$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$

$$-\eta = 1$$



• Iterated localized optimization with quadratic approximations

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^{T}$$

#### **Issues: 1. The Hessian**

• Normalized update rule

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E(\mathbf{w}^{(k)})^{-1} \nabla_{\mathbf{w}} E(\mathbf{w}^{(k)})^T$$

- For complex models such as neural networks, with a very large number of parameters, the Hessian  $H_E(w^{(k)})$  is extremely difficult to compute
  - For a network with only 100,000 parameters, the Hessian will have 10<sup>10</sup> cross-derivative terms
  - And its even harder to invert, since it will be enormous

#### **Issues: 1. The Hessian**



- For non-convex functions, the Hessian may not be positive semi-definite, in which case the algorithm can *diverge*
  - Goes away from, rather than towards the minimum

#### **Issues: 1. The Hessian**



- For non-convex functions, the Hessian may not be positive semi-definite, in which case the algorithm can *diverge*
  - Goes away from, rather than towards the minimum
  - Now requires additional checks to avoid movement in directions corresponding to –ve Eigenvalues of the Hessian

#### Issues: 1 – contd.

- A great many approaches have been proposed in the literature to *approximate* the Hessian in a number of ways and improve its positive definiteness
  - Boyden-Fletcher-Goldfarb-Shanno (BFGS)
    - And "low-memory" BFGS (L-BFGS)
    - Estimate Hessian from finite differences
  - Levenberg-Marquardt
    - Estimate Hessian from Jacobians
    - Diagonal load it to ensure positive definiteness
  - Other "Quasi-newton" methods
- Hessian estimates may even be *local* to a set of variables
- Not particularly popular anymore for large neural networks..

## **Issues: 2.** The learning rate



 Much of the analysis we just saw was based on trying to ensure that the step size was not so large as to cause divergence within a convex region

 $-\eta < 2\eta_{opt}$ 



• For complex models such as neural networks the loss function is often not convex

- Having  $\eta > 2\eta_{opt}$  can actually help escape local optima

• However *always* having  $\eta > 2\eta_{opt}$  will ensure that you never ever actually find a solution



- Start with a large learning rate
  - Greater than 2 (assuming Hessian normalization)
  - Gradually reduce it with iterations

# **Decaying learning rate**

• Typical decay schedules

– Linear decay: 
$$\eta_k = \frac{\eta_0}{k+1}$$

- Quadratic decay: 
$$\eta_k = \frac{\eta_0}{(k+1)^2}$$

– Exponential decay: 
$$\eta_k = \eta_0 e^{-\beta k}$$
, where  $\beta > 0$ 

- A common approach (for nnets):
  - 1. Train with a fixed learning rate  $\eta$  until loss (or performance on a held-out data set) stagnates
  - 2.  $\eta \leftarrow \alpha \eta$ , where  $\alpha < 1$  (typically 0.1)
  - 3. Return to step 1 and continue training from where we left off

## **Story so far : Convergence**

- Gradient descent can miss obvious answers
  - And this may be a good thing
- Convergence issues abound
  - The loss surface has many saddle points
    - Although, perhaps, not so many bad local minima
    - Gradient descent can stagnate on saddle points
  - Vanilla gradient descent may not converge, or may converge toooooo slowly
    - The optimal learning rate for one component may be too high or too low for others



Mark all true statements

- Step sizes that are greater than twice the inverse of the second derivative can cause gradient descent to diverge
- This is always a bad thing
- Gradient descent will not converge without decaying learning rates



Mark all true statements

- Step sizes that are greater than twice the inverse of the second derivative can cause gradient descent to diverge (true)
- This is always a bad thing
- Gradient descent will not converge without decaying learning rates

#### Story so far : Second-order methods

- Second-order methods "normalize" the variation along the components to mitigate the problem of different optimal learning rates for different components
  - But this requires computation of inverses of secondorder derivative matrices
  - Computationally infeasible
  - Not stable in non-convex regions of the loss surface
  - Approximate methods address these issues, but simpler solutions may be better
# **Story so far : Learning rate**

- Divergence-causing learning rates may not be a bad thing
  - Particularly for ugly loss functions
- Decaying learning rates provide good compromise between escaping poor local minima and convergence

• Many of the convergence issues arise because we force the same learning rate on all parameters

# Lets take a step back



$$\mathbf{w}^{(k+1)} \leftarrow \mathbf{w}^{(k)} - \eta (\nabla_{\mathbf{w}} E)^T$$

$$w_i^{(k+1)} = w_i^{(k)} - \frac{\eta}{\eta} \frac{dE\left(w_i^{(k)}\right)}{dw}$$

 Problems arise because of requiring a fixed step size across all dimensions

Because step are "tied" to the gradient

• Let's try releasing this requirement

# **Derivative-***inspired* algorithms

• Algorithms that use derivative information for trends, but do not follow them absolutely

- Rprop
- Quick prop

# RProp

- *Resilient* propagation
- Simple algorithm, to be followed *independently* for each component
  - I.e. steps in different directions are not coupled
- At each time
  - If the derivative at the current location recommends continuing in the same direction as before (i.e. has not changed sign from earlier):
    - *increase* the step, and continue in the same direction
  - If the derivative has changed sign (i.e. we've overshot a minimum)
    - *reduce* the step and reverse direction



- Select an initial value  $\widehat{w}$  and compute the derivative
  - Take an initial step  $\Delta w$  against the derivative
    - In the direction that reduces the function

$$-\Delta w = sign\left(\frac{dE(\widehat{w})}{dw}\right)\Delta w$$
$$-\widehat{w} = \widehat{w} - \Delta w$$



- Compute the derivative in the new location
  - If the derivative has not changed sign from the previous location, increase the step size and take a longer step

$$\alpha > 1$$
 •  $\Delta w = \alpha \Delta w$ 

•  $\widehat{w} = \widehat{w} - \Delta w$ 



- Compute the derivative in the new location
  - If the derivative has not changed sign from the previous location, increase the step size and take a step

$$\alpha > 1$$
 •  $\Delta w = \alpha \Delta w$ 

•  $\widehat{w} = \widehat{w} - \Delta w$ 



- Compute the derivative in the new location
  - If the derivative has changed sign



- Compute the derivative in the new location
  - If the derivative has changed sign
  - Return to the previous location
    - $\widehat{w} = \widehat{w} + \Delta w$



- Compute the derivative in the new location
  - If the derivative has changed sign
  - Return to the previous location
    - $\widehat{w} = \widehat{w} + \Delta w$
    - Shrink the step

B < 1

•  $\Delta w = \beta \Delta w$ 



- Compute the derivative in the new location
  - If the derivative has changed sign
  - Return to the previous location
    - $\widehat{w} = \widehat{w} + \Delta w$
    - Shrink the step

β<1

- $\Delta w = \beta \Delta w$
- Take the smaller step forward
  - $\widehat{w} = \widehat{w} \Delta w$

# **Rprop (simplified)**

- Set  $\alpha = 1.2, \beta = 0.5$
- For each layer *l*, for each *i*, *j*:
  - Initialize  $w_{l,i,j}$ ,  $\Delta w_{l,i,j} > 0$ ,

$$- prevD(l,i,j) = \frac{dLoss(w_{l,i,j})}{dw_{l,i,j}}$$

$$- \Delta w_{l,i,j} = \operatorname{sign}(prevD(l,i,j))\Delta w_{l,i,j}$$

- While not converged:

• 
$$w_{l,i,j} = w_{l,i,j} - \Delta w_{l,i,j}$$

• 
$$D(l, i, j) = \frac{dLoss(w_{l,i,j})}{dw_{l,i,j}}$$

- If sign(prevD(l,i,j)) == sign(D(l,i,j)):
  - $-\Delta w_{l,i,j} = \min(\alpha \Delta w_{l,i,j}, \Delta_{max}) \checkmark$
  - prevD(l, i, j) = D(l, i, j)
- else:

$$- w_{l,i,j} = w_{l,i,j} + \Delta w_{l,i,j}$$

 $- \Delta w_{l,i,j} = \max(\beta \Delta w_{l,i,j}, \Delta_{min}) \checkmark$ 

#### Ceiling and floor on step

# **Rprop (simplified)**

- Set  $\alpha = 1.2, \beta = 0.5$
- For each layer *l*, for each *i*, *j*:
  - Initialize  $w_{l,i,j}$ ,  $\Delta w_{l,i,j} > 0$ ,

 $- prevD(l, i, j) = \frac{dLoss(w_{l,i,j})}{dw_{l,i,j}}$ 

- $-\Delta w_{l,i,j} = \operatorname{sign}(prevD(l,i,j))\Delta w_{l,i,j}$
- While not converged:

• 
$$w_{l,i,j} = w_{l,i,j} - \Delta w_{l,i,j}$$

• 
$$D(l, i, j) = \frac{d Los(w_{l,i,j})}{d w_{l,i,j}}$$

• If  $\operatorname{sign}(\operatorname{prev} D(l, i, j)) == \operatorname{sign}(D(l, i, j))$ :

$$-\Delta w_{l,i,j} = \alpha \Delta w_{l,i,j}$$

- prevD(l, i, j) = D(l, i, j)
- else:

$$- w_{l,i,j} = w_{l,i,j} + \Delta w_{l,i,j}$$

$$- \Delta w_{l,i,j} = \beta \Delta w_{l,i,j}$$

Obtained via backprop

Note: Different parameters updated independently

# RProp

- A remarkably simple first-order algorithm, that is frequently much more efficient than gradient descent.
  - And can even be competitive against some of the more advanced second-order methods
- Only makes minimal assumptions about the loss function
  - No convexity assumption



The derivative of the loss w.r.t a parameter w, computed at the current estimate is positive. After taking a step (updating the parameter by a increment dw) the sign of the derivative becomes negative. Mark all true statements

- Rprop will revert to the earlier estimate and take a smaller step
- Rprop will change direction and begin taking steps in the opposite direction



The derivative of the loss w.r.t a parameter w, computed at the current estimate is positive. After taking a step (updating the parameter by a increment dw) the sign of the derivative becomes negative. Mark all true statements

- Rprop will revert to the earlier estimate and take a smaller step (true)
- Rprop will change direction and begin taking steps in the opposite direction

# QuickProp



• Quickprop employs the Newton updates with two modifications

$$\mathbf{w}^{(k+1)} = \mathbf{w}^{(k)} - \eta H_E \left( \mathbf{w}^{(k)} \right)^{-1} \nabla_{\mathbf{w}} E \left( \mathbf{w}^{(k)} \right)^T$$

• But with two modifications

# **QuickProp: Modification 1**



- It treats each dimension independently
- For i = 1: N

$$w_{i}^{k+1} = w_{i}^{k} - E''\left(w_{i}^{k}|w_{j}^{k}, j \neq i\right)^{-1}E'\left(w_{i}^{k}|w_{j}^{k}, j \neq i\right)$$

• This eliminates the need to compute and invert expensive Hessians

# **QuickProp: Modification 2**



- It approximates the second derivative through finite differences
- For i = 1: N

$$w_{i}^{k+1} = w_{i}^{k} - D(w_{i}^{k}, w_{i}^{k-1})^{-1} E'(w_{i}^{k}|w_{j}^{k}, j \neq i)$$

• This eliminates the need to compute expensive double derivatives

# QuickProp

$$w^{(k+1)} = w^{(k)} - \left(\frac{E'(w^{(k)}) - E'(w^{(k-1)})}{\Delta w^{(k-1)}}\right)^{-1} E'(w^{(k)})$$

Finite-difference approximation to double derivative obtained assuming a quadratic E()

- Updates are independent for every parameter
- For every layer l, for every connection from node i in the (l 1)<sup>th</sup> layer to node j in the l<sup>th</sup> layer:

$$\Delta w_{l,ij}^{(k)} = \frac{\Delta w_{l,ij}^{(k-1)}}{Err'\left(w_{l,ij}^{(k)}\right) - Err'\left(w_{l,ij}^{(k-1)}\right)} Err'\left(w_{l,ij}^{(k)}\right)$$

$$w_{l,ij}^{(k+1)} = w_{l,ij}^{(k)} - \Delta w_{l,ij}^{(k)}$$

# QuickProp

$$w^{(k+1)} = w^{(k)} - \left(\frac{E'(w^{(k)}) - E'(w^{(k-1)})}{\Delta w^{(k-1)}}\right)^{-1} E'(w^{(k)})$$

Finite-difference approximation to double derivative obtained assuming a quadratic E()

- Updates are independent for every parameter
- For every layer l, for every connection from node i in the (l 1)<sup>th</sup> layer to node j in the l<sup>th</sup> layer:

$$\Delta w_{l,ij}^{(k)} = \frac{\Delta w_{l,ij}^{(k-1)}}{Err'\left(w_{l,ij}^{(k)}\right) - Err'\left(w_{l,ij}^{(k-1)}\right)} \underbrace{Err'\left(w_{l,ij}^{(k)}\right)}_{k,ij} \underbrace{Err$$

# Quickprop

- Employs Newton updates with empirically derived derivatives
- Prone to some instability for non-convex objective functions

• But is still one of the fastest training algorithms for many problems

# **Story so far : Convergence**

- Gradient descent can miss obvious answers
  - And this may be a good thing
- Vanilla gradient descent may be too slow or unstable due to the differences between the dimensions
- Second order methods can normalize the variation across dimensions, but are complex
- Adaptive or decaying learning rates can improve convergence
- Methods that decouple the dimensions can improve convergence

# A closer look at the convergence problem



• With dimension-independent learning rates, the solution will converge smoothly in some directions, but oscillate or diverge in others

# A closer look at the convergence problem



- With dimension-independent learning rates, the solution will converge smoothly in some directions, but oscillate or diverge in others
- Proposal:
  - Keep track of oscillations
  - Emphasize steps in directions that converge smoothly
  - Shrink steps in directions that bounce around..

# The momentum methods

- Maintain a running average of all past steps
  - In directions in which the convergence is smooth, the average will have a large value
  - In directions in which the estimate swings, the positive and negative swings will cancel out in the average
- Update with the running average, rather than the current gradient





 The momentum method maintains a running average of all gradients until the *current* step

$$\Delta W^{(k)} = \beta \Delta W^{(k-1)} - \eta \nabla_W Loss \left( W^{(k-1)} \right)^{\mathsf{T}}$$
$$W^{(k)} = W^{(k-1)} + \Delta W^{(k)}$$

- Typical  $\beta$  value is 0.9
- The running average steps
  - Get longer in directions where gradient retains the same sign
  - Become shorter in directions where the sign keeps flipping

# **Training by gradient descent**

- Initialize all weights  $W_1, W_2, ..., W_K$
- Do:
  - For all *i*, *j*, *k*, initialize  $\nabla_{W_k} Loss = 0$
  - For all t = 1:T
    - For every layer k:
      - Compute  $\nabla_{W_k} Div(Y_t, d_t)$
      - Compute  $\nabla_{W_k} Loss += \frac{1}{T} \nabla_{W_k} Div(Y_t, d_t)$

- For every layer k:

 $W_k = W_k - \eta (\nabla_{W_k} Loss)^T$ 

Until Loss has converged

# **Training with momentum**

- Initialize all weights  $W_1, W_2, ..., W_K$
- Do:
  - For all layers k, initialize  $\nabla_{W_k} Loss = 0$ ,  $\Delta W_k = 0$
  - For all t = 1:T
    - For every layer k:
      - Compute gradient  $\nabla_{W_k} Div(Y_t, d_t)$
      - $-\nabla_{W_k}Loss += \frac{1}{T}\nabla_{W_k}\mathbf{Div}(Y_t, d_t)$
  - For every layer k

 $\Delta W_k = \beta \Delta W_k - \eta (\nabla_{W_k} Loss)^T$  $W_k = W_k + \Delta W_k$ 

Until Loss has converged



- The momentum method  $\Delta W^{(k)} = \beta \Delta W^{(k-1)} - \eta \nabla_W Loss(W^{(k-1)})^T$
- At any iteration, to compute the current step:



- The momentum method  $\Delta W^{(k)} = \beta \Delta W^{(k-1)} - \eta \nabla_W Loss(W^{(k-1)})^T$
- At any iteration, to compute the current step:
  First computes the gradient step at the current location



- The momentum method  $\Delta W^{(k)} = \beta \Delta W^{(k-1)} - \eta \nabla_W Loss(W^{(k-1)})^T$
- At any iteration, to compute the current step:
  - First computes the gradient step at the current location
  - Then adds in the scaled *previous* step
    - Which is actually a running average



• The momentum method

$$\Delta W^{(k)} = \beta \Delta W^{(k-1)} - \eta \nabla_W Loss(W^{(k-1)})^T$$

- At any iteration, to compute the current step:
  - First computes the gradient step at the current location
  - Then adds in the scaled *previous* step
    - Which is actually a running average
  - To get the final step



- Momentum update steps are actually computed in two stages
  - First: We take a step against the gradient at the current location
  - Second: Then we add a scaled version of the previous step
- The procedure can be made more optimal by reversing the order of operations..

### **Nestorov's Accelerated Gradient**



- Change the order of operations
- At any iteration, to compute the current step:

### **Nestorov's Accelerated Gradient**



- Change the order of operations
- At any iteration, to compute the current step:

First extend the previous step


- Change the order of operations
- At any iteration, to compute the current step:
  - First extend the previous step
  - Then compute the gradient step at the resultant position



- Change the order of operations
- At any iteration, to compute the current step:
  - First extend the previous step
  - Then compute the gradient step at the resultant position
  - Add the two to obtain the final step



• Nestorov's method  $\Delta W^{(k)} = \beta \Delta W^{(k-1)} - \eta \nabla_W Loss (W^{(k-1)} + \beta \Delta W^{(k-1)})^T$   $W^{(k)} = W^{(k-1)} + \Delta W^{(k)}$ 



- Comparison with momentum (example from Hinton)
- Converges much faster

## **Training with Nestorov**

- Initialize all weights **W**<sub>1</sub>, **W**<sub>2</sub>, ..., **W**<sub>K</sub>
- Do:
  - For all layers k, initialize  $\nabla_{W_k} Loss = 0$ ,  $\Delta W_k = 0$
  - For every layer k

 $W_k = W_k + \beta \Delta W_k$ 

- For all t = 1:T
  - For every layer k:
    - Compute gradient  $\nabla_{W_k} Div(Y_t, d_t)$

$$- \nabla_{W_k} Loss += \frac{1}{T} \nabla_{W_k} Div(Y_t, d_t)$$

For every layer k

$$W_{k} = W_{k} - \eta (\nabla_{W_{k}} Loss)^{T}$$
$$\Delta W_{k} = \beta \Delta W_{k} - \eta (\nabla_{W_{k}} Loss)^{T}$$

• Until *Loss* has converged

# Momentum and trend-based methods..

• We will return to this topic again, very soon..



On a flat surface of constant slope momentum methods will converge faster than vanilla gradient descent, true or false

- True
- False



On a flat surface of constant slope momentum methods will converge faster than vanilla gradient descent, true or false

- True
- False (correct) momentum only changes step size

## Story so far

- Gradient descent can miss obvious answers
  - And this may be a good thing
- Vanilla gradient descent may be too slow or unstable due to the differences between the dimensions
- Second order methods can normalize the variation across dimensions, but are complex
- Adaptive or decaying learning rates can improve convergence
- Methods that decouple the dimensions can improve convergence
- Momentum methods which emphasize directions of steady improvement are demonstrably superior to other methods

## Coming up

- Incremental updates
- Revisiting "trend" algorithms
- Generalization
- Tricks of the trade
  - Divergences..
  - Activations
  - Normalizations